

Laboratory Investigation Report

Patient Name : Mrs. Priti Kumari Yogendra Rai Centre : 2061 - Dr. Bnh, Dept. Of Laboratory Medicine

 Age/Gender
 : 33 Y 9 M 6 D /F
 OP/IP No/UHID
 : OP/MAXHOME/

 MaxID/Lab ID
 : NSSH.1237848/1816012520246
 Collection Date/Time
 : 26/Jan/2025 12:25PM

 Ref Doctor
 : Dr.Rishabh Nanavati
 Reporting Date/Time
 : 27/Jan/2025 03:32PM

Beta-2 Glycoprotein 1, IgM, Serum

Beta-2 Glycoprotein 1, IgM 2.9 U/mL < 7

FEIA

Result In U/ML	Remarks
Negative	< 7
Equivocal	7 - 10
Positive	> 10

Interpretation:

Detection of Beta-2 Glycoprotein antibodies are indicative of risk for thrombosis in autoimmune diseases.

Beta-2 Glycoprotein I IgM is intended for the in vitro quantitative measurement of IgM antibodies directed to β 2-Glycoprotein I in human serum and plasma to aid in the diagnosis of antiphospholipid syndrome (APS) and to evaluate the thrombotic risk in patients with systemic lupus erythematosus (SLE). A definitive clinical diagnosis should not be based on the results of a single diagnostic method, but should only be made after all clinical and laboratory findings have been evaluated. Rheumatoid factor (RF) can interfere with the determination of IgM anti- β 2-Glycoprotein I antibodies.





Laboratory Investigation Report

Patient Name : Mrs. Priti Kumari Yogendra Rai Centre : 2061 - Dr. Bnh, Dept. Of Laboratory Medicine

Age/Gender :33 Y 9 M 6 D /F OP/IP No/UHID : OP//MAXHOME/ : NSSH.1237848/1816012520246 MaxID/Lab ID Collection Date/Time: 26/Jan/2025 12:25PM Ref Doctor : Dr.Rishabh Nanavati Reporting Date/Time: 27/Jan/2025 03:32PM

Serology Special Bio Ref Interval **Test Name** Result Unit

Beta-2 Glycoprotein 1, IgG, Serum

Beta-2 Glycoprotein 1, IgG 0.6 U/mL < 7

Result In U/ML Remarks Negative < 7 Equivocal 7 - 10 > 10 Positive

Interpretation:

Detection of Beta-2 Glycoprotein antibodies are indicative of risk for thrombosis in autoimmune diseases.

Beta-2 Glycoprotein I IgG is intended for the in vitro quantitative measurement of IgG antibodies directed to β2-Glycoprotein I in human serum and plasma to aid in the diagnosis of antiphospholipid syndrome (APS) and to evaluate the thrombotic risk in patients with systemic lupus erythematosus (SLE). A definitive clinical diagnosis should not be based on the results of a single diagnostic method, but should only be made after all clinical and laboratory findings have been evaluated.

Kindly correlate with clinical findings

*** End Of Report ***

Dr.Poonam.S. Das, M.D. Principal Director-Max Lab & Blood Bank Services

Director, Microbiology & Molecular

Consultant Microbiology

Dr Nidhi Malik, MD Consultant Microbiology

Page 2 of 2

