



### Laboratory Investigation Report

Patient Name	Centre
Age/Gender	OP/IP No/UHID
MaxID/Lab ID	Collection Date/Time
Ref Doctor	Reporting Date/Time

Test Name	Result	Unit	Bio Ref Interval
-----------	--------	------	------------------

**HIV P24 Antigen (Screening)\*, Serum**

CLIA			
HIV P24 CMIA	Non Reactive		Non Reactive

**Comment Interpretation -**

p24 antigen can generally be detected in serum within next two weeks following exposure to HIV-1. Its presence reflects multiplication of the virus and hence provides an early diagnosis of the disease at the pre-sero conversion stage. In children from HIV-1 positive mothers, the presence of p 24 antigen at birth indicates early and severe forms of the disease. All p24 antigen reports should be correlated with HIV antibody and / or HIV viral load results.

Kindly correlate with clinical findings

\*\*\* End Of Report \*\*\*

  
**Dr. Poonam S. Das, M.D**  
 Principal Director  
 Max Lab & Blood Bank Services

  
**Dr. Bansidhar Tarai, M.D**  
 Associate Director  
 Microbiology & Molecular Diagnostics

  
**Dr. Sonu Kumari Agrawal, MD**  
 Associate Consultant  
 Microbiology